

HYPOSPADIA IN SMALL RUMINANTS: A CASE REPORT

Kalin Hristov, Aleksandar Stoimenov*

University of Forestry, Faculty of Veterinary Medicine, Sofia, Bulgaria

*e-mail: a.stoimenov@ltu.bg

ABSTRACT

Hypospadias is a rare congenital malformation of the urethra reported in dogs, sheep, goats, cattle, rats, nonhuman primates and humans. Affected animals may have other congenital or developmental anomalies such as cryptorchidism, which is to be the most common congenital anomaly associated with hypospadias.

In this clinical case is described condition of six weeks old male goat, weighing 10 kg, in Vrazhdebnna Training and Experimental Field Centre of the University of Forestry, with signs of urine scalding on the ventral perineum. Physical examination revealed aplasia of the penis and the urethra opened cranially to the scrotum. The diagnosis was made as hypospadias and penile aplasia.

Key words: hypospadias; penile aplasia; male goat

INTRODUCTION

Congenital malformations can result from defective genetics or environmental factors or a combination of both (Shukla et al., 2007). Pedigree analysis and breeding trials revealed that these anomalies are autosomal recessive diseases (Bryan et al., 1993). Congenital urinary tract anomalies in farm animals are rare. The most common anomalies are patent urachus, hypospadias, and renal agenesis (Temizosylu 2005). Hypospadias is imperfect closure of the external male urethra (Radostits et al., 2007). This congenital defect results from failure of fusion of the urogenital folds and incomplete formation of the penile urethra (Boothe, 2003). Hypospadias is accompanied by hypoplasia of the corpus cavernosum urethra (Fossum, 2002). Affected animals may have other congenital or developmental anomalies such as cryptorchidism, which is reported to be the most congenital anomaly associated with hypospadias. The aetiology of hypospadias is not well understood, it seems to be multifactorial and may be related to genetic, endocrinological, and environmental factors (Silver, 2000). In a survey of the occurrence of congenital anomalies in goats, the occurrence of congenital hypospadias was 0.066% (Al-Ani et al. 1998).

CASE PRESENTATION

The patient is a 6-week-old goat kid (*Capra hircus*), weighing 10 kg, grown at the Vrajzbenna Training and Experimental Center at the University of Forestry. The pregnancy of the mother was without complication and within the normal duration. Obstetric assistance is not applied at birth and only one kid is born, fully developed and vital.

After birth, the newborn started to suck milk from his mother within a few hours, and then goes on to an additional feeding. With age, animal keeper noticed persistence of urine in the area of the ventral perineum (Fig.1). During the clinical examination, it was established: BTT - 39.4 °C, pulse rate - 95 per minute, Respiratory rate - 27, preserved appetite and thirst. In the andrological examination, incomplete and improper development of the penile (penile hypoplasia) and prepuce, as well as open cranial to the scrotum urethra (Fig.2). The examination of the scrotum revealed the presence of two normally-developed testes for the age of the animal, with no change in their location - excluding cryptorchism (Fig.3).



Fig. 1 Presence of urine and pollution of the ventral perineum



Fig. 2 Penile hypoplasia and urethral hypospadias



Fig. 3 Normal development of the scrotum and testicular position



DISCUSSION

Hypospadias is a defect of the external genitalia characterized by an incomplete development of the prepuce with a ventral opening in the urethra at some point along the penis or the perineum due to lack of partial fusing of the urethral fold. In sheep has been regarded as a form of pseudohermaphroditism (Dennis and Leipold, 1979). In human, hypospadias is the second most common congenital abnormality after cryptorchidism (Pierik et al., 2002).

Hypospadias, urethral fissure at the ventral aspect of penile urethra, was mentioned in the available literatures (Sndak et al., 2010).

Three types of hypospadias are reported depending on the anatomical location of the urethral opening. The first (penile) form - the urethra opening could be proximal, distal, or in the mid shaft of the penis. The second form is scrotal in which the urethra opens between the halves of the divided scrotum and the third is the perineal in which the urethra opens in the perineum. Some authors classify hypospadias into 4 types: palanetic, penile, scrotal and perineal (Ladds, 1993).

In the present case the kid was born with this defect, which lead us to the conclusion that it is a congenital malformation. There are many reports concerning hypospadiac cases in ruminants associated with other congenital anomalies such as absence of tail, atresia ani, hermaphroditism, penile hypoplasia and cryptorchidism (Azari et al., 2010; Omidi et al., 2011; Bokhari, 2013; Saunders and Ladds, 1978). This is in accordance with the present case, in which we found hypospadias and hypoplasia of the penis and the prepuce. In this case, no treatment was given, but surgical reconstruction through permanent urethrostomy is the only possible option to treat hypospadias associated with penile hypoplasia. In case of surgical treatment of this condition, account should be taken of possible complication. Early complications of hypospadias repair include bleeding, hematoma, wound infection, wound dehiscence and urinary tract infection (Stokowski, 2004; Synder et al., 2005). Some late complications include urethrocutaneous fistula, urethral stricture, balanitis and urethrocele (Nuininga et al., 2005).

COMPETING INTERESTS

The authors declare that there is no conflict of interests regarding the publication of this paper.

ACKNOWLEDGEMENTS

The present study was initiated and implemented with the support of the National Science Program "Young Scientists and Postdoctoral Students". We express our gratitude for the concern for the young researchers, scientists and professors of Bulgaria

REFERENCES

1. Al-Ani F. K, Khamas W. A, Al-Qudah K. M, Al-Rawashdeh O., (1998) Occurrence of congenital anomalies in Shami breed goats: 211 cases investigated in 19 herds. Small Rumin Res 28(3):225–232
2. Azari O., Sakhaei E., Emadi L., (2010), Permanent Urethrostomy for Treatment of Caprine Hypospadias, American Journal of Animal and Veterinary Sciences 5 (2): 107-110,

3. Bokhari S. G., (2013). Hypospadias and urethral diverticulum in two goat kids: a case report, *The Journal of Animal & Plant Sciences*, 23(2): 675-677 ISSN: 1018-7081
4. Boothe, H.W., (2003). Penis, Prepuce and Scrotum. In: *Text Book of Small Animal Surgery*, Slatter, D. (Eds.), 3rd Edn., WB Saunders Company, Philadelphia, USA., ISBN: 0-7216-8607-9, pp: 153
5. Bryan L, Schmutz S, Hodges S. D, Snyder F. F (1993). Bovine b-mannosidosis: Pathologic and genetic findings in salers calves. *Vet. Pathol.* 30: 130 – 139.
6. Dennis, S.M., Leipold, H.W., (1979). Ovine congenital defects. *Vet. Bull.* 49, 233–239.
7. Fossum, T. W., 2002. Surgery of the Male Reproductive Tract. In: *Small Animal Surgery*, Fossum, T. W., C. S. Hedlund, D.A. Hulse, A.L. Johnson and H.B. Seim *et al.* (Eds.), 2nd Edn., Mosby, Inc., Elsevier Science Company, USA., ISBN: 0-32301238-8, pp: 662-663.
8. Ladds, P. W. (1993). Congenital Abnormalities of the Genitalia of Cattle Sheep, Goats, and Pigs. *The Veterinary Clinics of North America: Food Animal Practice*, 9, 127-144.
9. Nuininga, J.E., R.P.E. de Gier, R. Vetschuren and W.F. Feitz, (2005). Long-term outcome of different types of 1-stage hypospadias repairs. *J. Urol.*, 174: 1544-1548. PMID: 16148649
10. Omidi A., S. Monjezi, and A. Nabipour, (2011). Hypospadias, diverticulum, and agenesis in the penile shaft of a goat kid (*Capra hircus*), *Comparative Clinical Pathology*, vol. 20, no. 4, pp. 417–419,
11. Pierik, F.H., A. Burdorf, J.M.R. Nijman, R.E. Juttmann and S.M.P.F. de Munick Keizer. (2002). A high hypospadias rate in the Netherlands. *Hum. Reprod.*, 17: 1112-1115. PMID: 11925415
12. Radostits, O. M., C. C. Gay, K.W. Hinchcliff and P. D. Constable, (2007). *Veterinary Medicine*. 10th Ed., Saunders Co., London, ISBN: 0-7020-2777-4, pp: 573.
13. Saunders P.J, Ladds P. W., (1978). Congenital and developmental anomalies of the genitalia of slaughtered bulls, *Aust Vet J.* Jan, 54(1):10-3, doi.org/10.1111/j.1751-0813.1978.tb00261.x
14. Shukla S. P., Nema S. P., Pandey A. K., Garg U. K. (2007). Dystocia due to a bull dog calf in a she buffalo. *Buff. Bull.* 26: 104 – 105.
15. Silver, R. I., (2000). What is the etiology of hypospadias, a review of recent research. *Delaware Med. J.*, 72: 343-347. PMID: 10984977
16. Sndak, N., Sahn, T. and Brck, H. S. (2010). Urethral Dilatation, Ectopic Testis, Hypoplasia Penis, and Phimosis in a Kilis Goat Kid," *Kafkas universitesi veteriner fakultesi dergisi*, 16, 147-150.
17. Stokowski, L. A., (2004). Hypospadias in neonate. *Adv.Neonatal Care*, 4: 463-472. PMID: 15368213
18. Synder, C. L., A. Evangelidis, G. Hasen, S. D. Peter and D. J. Ostlie. (2005). Management of complications after hypospadias repair. *Urology*, 65: 782-785. PMID: 15833528
19. Temizosyulu M. D. (2005) Penile urethral diverticulum in a kid. *Ankara Univ Vet Fak Derg* 52:185–187

